## Years of Potential Life Lost (YPLL) before Age 65, by Cause of Death, 2000-2009, by Race and Hispanic Ethnicity

White, non-Hispanic				
Rank	Cause	Average	Percent of	
		Annual YPLL	Total YPLL	
	All Causes	41,657		
1	Unintentional Injury	9,288	22.3%	
2	Malignant Neoplasms	8,882	21.3%	
3	Heart Disease	5,253	12.6%	
4	Suicide	3,185	7.6%	
5	Perinatal Period	2,461	5.9%	
	All Others	12,587	30.2%	

Black, non-Hispanic				
Rank	Cause	Average Annual YPLL	Percent of Total YPLL	
	All Causes	610		
1	Unintentional Injury	142	23.3%	
2	Perinatal Period	98	16.0%	
3	Heart Disease	70	11.4%	
4	Malignant Neoplasms	46	7.5%	
5	Congenital Anomalies	44	7.3%	
	All Others	211	34.6%	

Native American, non-Hispanic			
Double	Cause	Average	Percent of
Rank		Annual YPLL	Total YPLL
All Causes		443	
1	Unintentional Injury	111	25.0%
2	Heart Disease	62	13.9%
3	Malignant Neoplasms	61	13.8%
4	Suicide	33	7.5%
5	Diabetes Mellitus	21	4.7%
	All Others	155	35.1%

Asian/Pacific Islander, non-Hispanic			
Rank	Cause	Average	Percent of
		Annual YPLL	Total YPLL
	All Causes	247	
1	Malignant Neoplasms	57	22.9%
2	Unintentional Injury	55	22.0%
3	Congenital Anomalies	33	13.1%
4	Perinatal Period	26	10.5%
5	Heart Disease	17	6.7%
	All Others	61	24.7%

Other, non-Hispanic			
Rank	Cause		Percent of
			Total YPLL
	All Causes	690	
1	Unintentional Injury	165	23.9%
2	Malignant Neoplasms	118	17.1%
3	Heart Disease	78	11.3%
4	Congenital Anomalies	50	7.2%
5	Suicide	50	7.2%
	All Others	229	33.2%

	His	panic	
Rank	Cause	Average	Percent of
		Annual YPLL	Total YPLL
	All Causes	272	
1	Unintentional Injury	71	26.2%
2	Malignant Neoplasms	27	9.9%
3	Suicide	26	9.7%
4	Perinatal Period	20	7.2%
5	Congenital Anomalies	18	6.6%
	All Others	110	40.5%

Data accessed on WISQARS, US CDC. http://www.cdc.gov/injury/wisqars/years\_potential.html .

Because numbers are not controlled for population size, YPLL can not be compared across different population groups; only the order of causes can be compared.